

Self sustained traversable wormholes and the equation of state

Remo Garattini*

*Università degli Studi di Bergamo, Facoltà di Ingegneria,
Viale Marconi 5, 24044 Dalmine (Bergamo) ITALY.*

We compute the graviton one loop contribution to a classical energy in a *traversable* wormhole background. The form of the shape function considered is obtained by the equation of state $p = \omega\rho$. We investigate the size of the wormhole as a function of the parameter ω . The investigation is evaluated by means of a variational approach with Gaussian trial wave functionals. A zeta function regularization is involved to handle with divergences. A renormalization procedure is introduced and the finite one loop energy is considered as a *self-consistent* source for the traversable wormhole. The case of the phantom region is briefly discussed.

I. INTRODUCTION

The discovery that our universe is undergoing an accelerated expansion[1] leads to reexamine the Friedmann-Robertson-Walker equation

$$\frac{\ddot{a}}{a} = -\frac{4\pi}{3}(\rho + 3p), \quad (1)$$

to explain why the scale factor obeys $\ddot{a} > 0$. Indeed, it is evident from the previous formula, that a sort of *dark energy* is needed to cause a negative pressure with equation of state

$$p = \omega\rho. \quad (2)$$

A value of $\omega < -1/3$ is required for the accelerated expansion, while $\omega = -1$ corresponds to a cosmological constant. A specific form of dark energy, denoted *phantom energy* has also been proposed with the property of having $\omega < -1$. It is interesting to note that the phantom energy violates the null energy condition, $p + \rho < 0$, necessary ingredient to sustain the traversability of wormholes. A wormhole can be represented by two asymptotically flat regions joined by a bridge. To exist, it must satisfy the Einstein field equations: one example is represented by the Schwarzschild solution. One of the prerogatives of a wormhole is its ability to connect two distant points in space-time. In this amazing perspective, it is immediate to recognize the possibility of traveling crossing wormholes as a short-cut in space and time. Unfortunately, although there is no direct evidence, a Schwarzschild wormhole does not possess this property. It is for this reason that in a pioneering work Morris and Thorne[2] and subsequently Morris, Thorne and Yurtsever[3] studied a class of wormholes termed “*traversable*”. Unfortunately, the traversability is accompanied by unavoidable violations of null energy conditions, namely, the matter threading the wormhole’s throat has to be “*exotic*”. It is clear that the existence of dark and phantom energy supports the class of exotic matter. In this direction, Lobo[4], Kuhfittig[5] and Sushkov[6] have considered the possibility of sustaining the wormhole traversability with the help of phantom energy. In a previous work, we explored the possibility that a wormhole can be sustained by its own quantum fluctuations[7]. In practice, it is the graviton propagating on the wormhole background that plays the role of the “*exotic*” matter. This has not to appear as a surprise, because the computation involved, namely the one loop contribution of the graviton to the total energy, is quite similar to compute the Casimir energy on a fixed background. It is known that, for different physical systems, Casimir energy is negative and this is exactly one of the features that the exotic matter should possess. In particular, we conjectured that quantum fluctuations can support the traversability as effective source of the semiclassical Einstein’s equations. However in Ref.[7], we limited the analysis in the region where the equation of state(2) assumes the particular value $\omega = 1$. In this paper, we will consider $\omega \in (0, +\infty)$ [8], although the semiclassical approach can be judged suspicious because of the suspected validity of semiclassical methods¹ at the Planck scale[9]. The rest of the paper is structured as follows, in section II we define the effective Einstein equations, in section III we introduce the traversable wormhole metric, in section IV we give some of the basic rules to perform the functional integration and we define the Hamiltonian approximated up to second order, in section V we study the spectrum of the spin-two operator acting on transverse traceless tensors,

*Electronic address: Remo.Garattini@unibg.it

¹ To this purpose, see also paper of Hochberg, Popov and Sushkov[10] and the paper of Khusnutdinov and Sushkov[11].

in section VI we regularize and renormalize the one loop energy contribution and we speculate about self-consistency of the result. We summarize and conclude in section VII.

II. THE EFFECTIVE EINSTEIN EQUATIONS

We begin with a look at the classical Einstein equations

$$G_{\mu\nu} = \kappa T_{\mu\nu}, \quad (3)$$

where $T_{\mu\nu}$ is the stress-energy tensor, $G_{\mu\nu}$ is the Einstein tensor and $\kappa = 8\pi G$. Consider a separation of the metric into a background part, $\bar{g}_{\mu\nu}$, and a perturbation, $h_{\mu\nu}$,

$$g_{\mu\nu} = \bar{g}_{\mu\nu} + h_{\mu\nu}. \quad (4)$$

The Einstein tensor $G_{\mu\nu}$ can also be divided into a part describing the curvature due to the background geometry and that due to the perturbation,

$$G_{\mu\nu}(g_{\alpha\beta}) = G_{\mu\nu}(\bar{g}_{\alpha\beta}) + \Delta G_{\mu\nu}(\bar{g}_{\alpha\beta}, h_{\alpha\beta}), \quad (5)$$

where, in principle $\Delta G_{\mu\nu}(\bar{g}_{\alpha\beta}, h_{\alpha\beta})$ is a perturbation series in terms of $h_{\mu\nu}$. In the context of semiclassical gravity, Eq.(3) becomes

$$G_{\mu\nu} = \kappa \langle T_{\mu\nu} \rangle^{ren}, \quad (6)$$

where $\langle T_{\mu\nu} \rangle^{ren}$ is the renormalized expectation value of the stress-energy tensor operator of the quantized field. If the matter field source is absent, nothing prevents us from defining an effective stress-energy tensor for the fluctuations as²

$$\langle T_{\mu\nu} \rangle^{ren} = -\frac{1}{\kappa} \langle \Delta G_{\mu\nu}(\bar{g}_{\alpha\beta}, h_{\alpha\beta}) \rangle^{ren}. \quad (7)$$

From this point of view, the equation governing quantum fluctuations behaves as a backreaction equation. If we fix our attention to the energy component of the Einstein field equations, we need to introduce a time-like unit vector u^μ such that $u \cdot u = -1$. Then the semi-classical Einstein's equations (6) projected on the constant time hypersurface Σ become

$$G_{\mu\nu}(\bar{g}_{\alpha\beta}) u^\mu u^\nu = \kappa \langle T_{\mu\nu} u^\mu u^\nu \rangle^{ren} = -\langle \Delta G_{\mu\nu}(\bar{g}_{\alpha\beta}, h_{\alpha\beta}) u^\mu u^\nu \rangle^{ren}. \quad (8)$$

To further proceed, it is convenient to consider the associated tensor density and integrate over Σ . This leads to³

$$\frac{1}{2\kappa} \int_\Sigma d^3x \sqrt{^3\bar{g}} G_{\mu\nu}(\bar{g}_{\alpha\beta}) u^\mu u^\nu = - \int_\Sigma d^3x \mathcal{H}^{(0)} = -\frac{1}{2\kappa} \int_\Sigma d^3x \sqrt{^3\bar{g}} \langle \Delta G_{\mu\nu}(\bar{g}_{\alpha\beta}, h_{\alpha\beta}) u^\mu u^\nu \rangle^{ren}, \quad (9)$$

where

$$\mathcal{H}^{(0)} = \frac{2\kappa}{\sqrt{^3\bar{g}}} G_{ijkl} \pi^{ij} \pi^{kl} - \frac{1}{2\kappa} \sqrt{^3\bar{g}} R^{(3)} \quad (10)$$

is the background field super-hamiltonian and G_{ijkl} is the DeWitt super metric. Thus the fluctuations in the Einstein tensor are, in this context, the fluctuations of the hamiltonian. To compute the expectation value of the perturbed Einstein tensor in the transverse-traceless sector, we use a variational procedure with gaussian wave functionals. In practice, the right hand side of Eq.(9) will be obtained by expanding

$$E_{wormhole} = \frac{\langle \Psi | H_\Sigma | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \frac{\langle \Psi | H_\Sigma^{(0)} + H_\Sigma^{(1)} + H_\Sigma^{(2)} + \dots | \Psi \rangle}{\langle \Psi | \Psi \rangle} \quad (11)$$

² Note that our approach is very close to the gravitational *geon* considered by Anderson and Brill[13]. The relevant difference is in the averaging procedure.

³ Details on sign conventions and decomposition of the Einstein tensor can be found in Apeendix B

and retaining only quantum fluctuations contributing to the effective stress energy tensor. $H_\Sigma^{(i)}$ represents the hamiltonian approximated to the i^{th} order in h_{ij} and Ψ is a *trial wave functional* of the gaussian form. Then Eq.(9) becomes

$$H_\Sigma^{(0)} = \int_\Sigma d^3x \mathcal{H}^{(0)} = - \frac{\langle \Psi | H_\Sigma^{(1)} + H_\Sigma^{(2)} + \dots | \Psi \rangle}{\langle \Psi | \Psi \rangle}. \quad (12)$$

The chosen background to compute the quantity contained in Eq.(9) will be that of a traversable wormhole.

III. EINSTEIN FIELD EQUATIONS AND THE TRAVERSABLE WORMHOLE METRIC

In Schwarzschild coordinates, the traversable wormhole metric can be cast into the form

$$ds^2 = -\exp(-2\phi(r)) dt^2 + \frac{dr^2}{1 - \frac{b(r)}{r}} + r^2 [d\theta^2 + \sin^2\theta d\varphi^2]. \quad (13)$$

where $\phi(r)$ is called the redshift function, while $b(r)$ is called the shape function. Using the Einstein field equation

$$G_{\mu\nu} = \kappa T_{\mu\nu}, \quad (14)$$

in an orthonormal reference frame, we obtain the following set of equations

$$\rho(r) = \frac{1}{8\pi G} \frac{b'}{r^2}, \quad (15)$$

$$p_r(r) = \frac{1}{8\pi G} \left[\frac{2}{r} \left(1 - \frac{b(r)}{r} \right) \phi' - \frac{b}{r^3} \right], \quad (16)$$

$$p_t(r) = \frac{1}{8\pi G} \left(1 - \frac{b(r)}{r} \right) \left[\phi'' + \phi' \left(\phi' + \frac{1}{r} \right) \right] - \frac{b'r - b}{2r^2} \left(\phi' + \frac{1}{r} \right), \quad (17)$$

in which $\rho(r)$ is the energy density, $p_r(r)$ is the radial pressure, and $p_t(r)$ is the lateral pressure. Using the conservation of the stress-energy tensor, in the same orthonormal reference frame, we get

$$p'_r = \frac{2}{r} (p_t - p_r) - (\rho + p_r) \phi'. \quad (18)$$

The Einstein equations can be rearranged to give

$$b' = 8\pi G \rho(r) r^2, \quad (19)$$

$$\phi' = \frac{b + 8\pi G p_r r^3}{2r^2 \left(1 - \frac{b(r)}{r} \right)}. \quad (20)$$

Now, we introduce the equation of state $p_r = \omega\rho$, and using Eq.(15), Eq.(20) becomes

$$\phi' = \frac{b + \omega b' r}{2r^2 \left(1 - \frac{b(r)}{r} \right)}. \quad (21)$$

The redshift function can be set to a constant with respect to the radial distance, if

$$b + \omega b' r = 0. \quad (22)$$

The integration of this simple equation leads to

$$b(r) = r_t \left(\frac{r_t}{r} \right)^{\frac{1}{\omega}}, \quad (23)$$

where we have used the condition $b(r_t) = r_t$. In this situation, the line element (13) becomes

$$ds^2 = -A dt^2 + \frac{dr^2}{1 - \left(\frac{r_t}{r}\right)^{1+\frac{1}{\omega}}} + r^2 [d\theta^2 + \sin^2 \theta d\varphi^2], \quad (24)$$

where A is a constant coming from $\phi' = 0$ which can be set to one without loss of generality. The parameter ω is restricted by the following conditions

$$b'(r_t) < 1; \quad \frac{b(r)}{r} \xrightarrow{r \rightarrow +\infty} 0 \quad \implies \quad \begin{cases} \omega > 0 \\ \omega < -1 \end{cases}. \quad (25)$$

Proper radial distance is related to the shape function by

$$l(r) = \pm \int_{r_t}^r \frac{dr'}{\sqrt{1 - \frac{b_{\pm}(r')}{r'}}} = \pm r_t \frac{2\omega}{\omega + 1} \sqrt{\rho^{(1+\frac{1}{\omega})} - 1} {}_2F_1 \left(\frac{1}{2}, \frac{1-\omega}{2\omega+2}; \frac{3}{2}; 1 - \rho^{(1+\frac{1}{\omega})} \right), \quad (26)$$

where the plus (minus) sign is related to the upper (lower) part of the wormhole or universe and where ${}_2F_1(a, b; c; x)$ is a hypergeometric function. Two coordinate patches are required, each one covering the range $[r_t, +\infty)$. Each patch covers one universe, and the two patches join at r_t , the throat of the wormhole defined by

$$r_t = \min \{r(l)\}. \quad (27)$$

When $\omega = 1$, we recover the special case where $b(r) = r_t^2/r$, then the line element simply becomes

$$ds^2 = -dt^2 + \frac{dr^2}{1 - \frac{r_t^2}{r^2}} + r^2 [d\theta^2 + \sin^2 \theta d\varphi^2] \quad (28)$$

and Eq.(26) simplifies into

$$l(r) = \pm \int_{r_t}^r \frac{dr'}{\sqrt{1 - \frac{r_t^2}{r'^2}}} = \pm \sqrt{r^2 - r_t^2} \quad \implies \quad r^2 = l^2 + r_t^2. \quad (29)$$

The new coordinate l covers the range $-\infty < l < +\infty$. The constant time hypersurface Σ is an Einstein-Rosen bridge with wormhole topology $S^2 \times R^1$. The Einstein-Rosen bridge defines a bifurcation surface dividing Σ in two parts denoted by Σ_+ and Σ_- . To concretely compute Eq.(12), we consider on the slice Σ deviations from the wormhole metric of the type

$$g_{ij} = \bar{g}_{ij} + h_{ij}, \quad (30)$$

where g_{ij} is extracted from the line element (24) whose form becomes

$$ds^2 = -dt^2 + g_{ij} dx^i dx^j. \quad (31)$$

IV. ENERGY DENSITY CALCULATION IN SCHRÖDINGER REPRESENTATION

In order to compute the quantity

$$- \int_{\Sigma} d^3x \sqrt{^3g} \langle \Delta G_{\mu\nu} (\bar{g}_{\alpha\beta}, h_{\alpha\beta}) u^{\mu} u^{\nu} \rangle^{ren}, \quad (32)$$

we consider the right hand side of Eq.(12). Since $H_{\Sigma}^{(1)}$ is linear in h_{ij} and h , the corresponding gaussian integral disappears and since

$$\frac{\sqrt{^3g}}{\kappa} G_{\mu\nu} (g_{\alpha\beta}) u^{\mu} u^{\nu} = -\mathcal{H}, \quad (33)$$

it is clear that the hamiltonian expansion in Eq.(12) does not coincide with the averaged expanded Einstein tensor of Eq.(32) because Eq.(33) involves a tensor density. Therefore, the correct setting is

$$\int_{\Sigma} d^3x \sqrt{^3g} \langle \Delta G_{\mu\nu} (\bar{g}_{\alpha\beta}, h_{\alpha\beta}) u^{\mu} u^{\nu} \rangle^{ren} = \int_{\Sigma} d^3x \sqrt{^3g} \frac{\langle \Psi | \mathcal{H}^{(2)} - \sqrt{^3g}^{(2)} \mathcal{H}^{(0)} | \Psi \rangle}{\langle \Psi | \Psi \rangle}, \quad (34)$$

where $\sqrt{^3g}^{(2)}$ is the second order expanded tensor density weight. Following the same procedure of Refs.[15, 16], the potential part of the right hand side of Eq.(34) becomes

$$\int_{\Sigma} d^3x \sqrt{g} \left[-\frac{1}{4} h \Delta h + \frac{1}{4} h^{li} \Delta h_{li} - \frac{1}{2} h^{ij} \nabla_l \nabla_i h_j^l + \frac{1}{2} h \nabla_l \nabla_i h^{li} - \frac{1}{2} h^{ij} R_{ia} h_j^a + \frac{1}{2} h R_{ij} h^{ij} + \frac{1}{4} R h^{ij} h_{ij} - \frac{1}{4} h R h \right]. \quad (35)$$

The term

$$\int_{\Sigma} d^3x \sqrt{g} \left[\frac{1}{4} R h^{ij} h_{ij} - \frac{1}{4} h R h \right], \quad (36)$$

makes the difference between the hamiltonian expansion and the Einstein tensor expansion. To explicitly make calculations, we need an orthogonal decomposition for both π_{ij} and h_{ij} to disentangle gauge modes from physical deformations. We define the inner product

$$\langle h, k \rangle := \int_{\Sigma} \sqrt{g} G^{ijkl} h_{ij}(x) k_{kl}(x) d^3x, \quad (37)$$

by means of the inverse WDW metric G_{ijkl} , to have a metric on the space of deformations, i.e. a quadratic form on the tangent space at h , with

$$G^{ijkl} = (g^{ik} g^{jl} + g^{il} g^{jk} - 2g^{ij} g^{kl}). \quad (38)$$

The inverse metric is defined on co-tangent space and it assumes the form

$$\langle p, q \rangle := \int_{\Sigma} \sqrt{g} G_{ijkl} p^{ij}(x) q^{kl}(x) d^3x, \quad (39)$$

so that

$$G^{ijnm} G_{nmkl} = \frac{1}{2} \left(\delta_k^i \delta_l^j + \delta_l^i \delta_k^j \right). \quad (40)$$

Note that in this scheme the “inverse metric” is actually the WDW metric defined on phase space. The desired decomposition on the tangent space of 3-metric deformations[17, 18] is:

$$h_{ij} = \frac{1}{3} h g_{ij} + (L\xi)_{ij} + h_{ij}^{\perp} \quad (41)$$

where the operator L maps ξ_i into symmetric tracefree tensors

$$(L\xi)_{ij} = \nabla_i \xi_j + \nabla_j \xi_i - \frac{2}{3} g_{ij} (\nabla \cdot \xi). \quad (42)$$

Thus the inner product between three-geometries becomes

$$\begin{aligned} \langle h, h \rangle &:= \int_{\Sigma} \sqrt{g} G^{ijkl} h_{ij}(x) h_{kl}(x) d^3x = \\ &\int_{\Sigma} \sqrt{g} \left[-\frac{2}{3} h^2 + (L\xi)^{ij} (L\xi)_{ij} + h^{ij\perp} h_{ij}^{\perp} \right]. \end{aligned} \quad (43)$$

With the orthogonal decomposition in hand we can define the trial wave functional

$$\Psi \{h_{ij}(\vec{x})\} = \mathcal{N} \exp \left\{ -\frac{1}{4l_p^2} \left[\langle hK^{-1}h \rangle_{x,y}^\perp + \langle (L\xi) K^{-1} (L\xi) \rangle_{x,y}^\parallel + \langle hK^{-1}h \rangle_{x,y}^{Trace} \right] \right\}, \quad (44)$$

where $\mathcal{N}$ is a normalization factor. We are interested in perturbations of the physical degrees of freedom. Thus we fix our attention only to the TT tensor sector reducing therefore the previous form into

$$\Psi \{h_{ij}(\vec{x})\} = \mathcal{N} \exp \left\{ -\frac{1}{4} \langle hK^{-1}h \rangle_{x,y}^\perp \right\}. \quad (45)$$

Therefore to calculate the energy density, we need to know the action of some basic operators on $\Psi[h_{ij}]$. The action of the operator h_{ij} on $|\Psi\rangle = \Psi[h_{ij}]$ is realized by

$$h_{ij}(x)|\Psi\rangle = h_{ij}(\vec{x})\Psi\{h_{ij}\}. \quad (46)$$

The action of the operator π_{ij} on $|\Psi\rangle$, in general, is

$$\pi_{ij}(x)|\Psi\rangle = -i \frac{\delta}{\delta h_{ij}(\vec{x})} \Psi\{h_{ij}\}. \quad (47)$$

The inner product is defined by the functional integration:

$$\langle \Psi_1 | \Psi_2 \rangle = \int [\mathcal{D}h_{ij}] \Psi_1^* \{h_{ij}\} \Psi_2 \{h_{kl}\}, \quad (48)$$

and by applying previous functional integration rules, we obtain the expression of the one-loop-like Hamiltonian form for TT (traceless and transverse) deformations

$$H_\Sigma^\perp = \frac{1}{4} \int_\Sigma d^3x \sqrt{g} G^{ijkl} \left[(16\pi G) K^{-1\perp}(x, x)_{ijkl} + \frac{1}{(16\pi G)} (\Delta_2)_j^a K^\perp(x, x)_{iakl} \right]. \quad (49)$$

The propagator $K^\perp(x, x)_{iakl}$ comes from a functional integration and it can be represented as

$$K^\perp(\vec{x}, \vec{y})_{iakl} := \sum_\tau \frac{h_{ia}^{(\tau)\perp}(\vec{x}) h_{kl}^{(\tau)\perp}(\vec{y})}{2\lambda(\tau)}, \quad (50)$$

where $h_{ia}^{(\tau)\perp}(\vec{x})$ are the eigenfunctions of Δ_2 , whose eigenvalues will be denoted with $\tilde{E}_i^2(\tau)$. τ denotes a complete set of indices and $\lambda(\tau)$ are a set of variational parameters to be determined by the minimization of Eq.(49). The expectation value of $H^\perp$ is easily obtained by inserting the form of the propagator into Eq.(49)

$$E(\lambda_i) = \frac{1}{4} \sum_\tau \sum_{i=1}^2 \left[(16\pi G) \lambda_i(\tau) + \frac{\tilde{E}_i^2(\tau)}{(16\pi G) \lambda_i(\tau)} \right]. \quad (51)$$

By minimizing with respect to the variational function $\lambda_i(\tau)$, we obtain the total one loop energy for TT tensors

$$E^{TT} = \frac{1}{4} \sum_\tau \left[\sqrt{\tilde{E}_1^2(\tau)} + \sqrt{\tilde{E}_2^2(\tau)} \right]. \quad (52)$$

The above expression makes sense only for $\tilde{E}_i^2(\tau) > 0$, $i = 1, 2$. The meaning of $\tilde{E}_i^2$ will be clarified in the next section. Coming back to Eq.(12), we observe that the value of the wormhole energy on the chosen background is⁴

$$\int_\Sigma d^3x \mathcal{H}^{(0)} = -\frac{1}{16\pi G} \int_\Sigma d^3x \sqrt{g} R^{(3)} = A(\omega) \frac{r_t}{G}, \quad \omega > -1 \quad (53)$$

⁴ Details of the calculation can be found in the Appendix A.

where

$$A(\omega) = \frac{1}{1+\omega} B\left(\frac{1}{2}, \frac{1}{1+\omega}\right) = \frac{\sqrt{\pi}}{(1+\omega)} \frac{\Gamma\left(\frac{1}{1+\omega}\right)}{\Gamma\left(\frac{3+\omega}{2+2\omega}\right)}. \quad (54)$$

$B(x, y)$ is the Beta function and $\Gamma(x)$ is the gamma function. Then the one loop the self-consistent equation for TT tensors becomes

$$A(\omega) \frac{r_t}{G} = -E^{TT}. \quad (55)$$

Note that for the special value of $\omega = 1$, we get

$$\frac{\pi r_t}{2G} = -E^{TT}, \quad (56)$$

in agreement with the result of Ref. [7]. Note also that the self-consistency on the hamiltonian as a reversed sign with respect to the energy component of the Einstein field equations. This means that an eventual stable point for the hamiltonian is an unstable point for the effective energy momentum tensor and vice versa.

V. THE TRANSVERSE TRACELESS (TT) SPIN 2 OPERATOR FOR THE TRAVERSABLE WORMHOLE AND THE W.K.B. APPROXIMATION

In this section, we evaluate the one loop energy expressed by Eq.(52). To this purpose, we begin with the operator describing gravitons propagating on the background (24). The Lichnerowicz operator in this particular metric is defined by

$$(\Delta_2 h^{TT})_i^j := -(\Delta_T h^{TT})_i^j + 2(Rh^{TT})_i^j - R(h^{TT})_i^j, \quad (57)$$

where the transverse-traceless (TT) tensor for the quantum fluctuation is obtained by the following decomposition

$$h_i^j = h_i^j - \frac{1}{3}\delta_i^j h + \frac{1}{3}\delta_i^j h = (h^T)_i^j + \frac{1}{3}\delta_i^j h. \quad (58)$$

This implies that $(h^T)_i^i = 0$. The transversality condition is applied on $(h^T)_i^j$ and becomes $\nabla_j (h^T)_i^j = 0$. Thus

$$-(\Delta_T h^{TT})_i^j = -\Delta_S (h^{TT})_i^j + \frac{6}{r^2} \left(1 - \frac{b(r)}{r}\right), \quad (59)$$

where Δ_S is the scalar curved Laplacian, whose form is

$$\Delta_S = \left(1 - \frac{b(r)}{r}\right) \frac{d^2}{dr^2} + \left(\frac{4r - b'(r)r - 3b(r)}{2r^2}\right) \frac{d}{dr} - \frac{L^2}{r^2} \quad (60)$$

and R_j^a is the mixed Ricci tensor whose components are:

$$R_i^a = \left\{ \frac{b'(r)}{r^2} - \frac{b(r)}{r^3}, \frac{b'(r)}{2r^2} + \frac{b(r)}{2r^3}, \frac{b'(r)}{2r^2} + \frac{b(r)}{2r^3} \right\}. \quad (61)$$

The scalar curvature is

$$R = R_i^j \delta_j^i = 2 \frac{b'(r)}{r^2} \quad (62)$$

We are therefore led to study the following eigenvalue equation

$$(\Delta_2 h^{TT})_i^j = \tilde{E}^2 h_i^j \quad (63)$$

where ω^2 is the eigenvalue of the corresponding equation. In doing so, we follow Regge and Wheeler in analyzing the equation as modes of definite frequency, angular momentum and parity[19]. In particular, our choice for the three-dimensional gravitational perturbation is represented by its even-parity form

$$h_{ij}^{even}(r, \vartheta, \phi) = \text{diag} \left[H(r) \left(1 - \frac{b(r)}{r} \right)^{-1}, r^2 K(r), r^2 \sin^2 \vartheta L(r) \right] Y_{lm}(\vartheta, \phi), \quad (64)$$

with

$$\begin{cases} H(r) = h_1^1(r) - \frac{1}{3}h(r) \\ K(r) = h_2^2(r) - \frac{1}{3}h(r) \\ L(r) = h_3^3(r) - \frac{1}{3}h(r) \end{cases}. \quad (65)$$

From the transversality condition we obtain $h_2^2(r) = h_3^3(r)$. Then $K(r) = L(r)$. For a generic value of the angular momentum L , representation (64) joined to Eq.(59) lead to the following system of PDE's

$$\begin{cases} \left(-\Delta_l + 2 \left(\frac{b'(r)}{r^2} - \frac{b(r)}{r^3} - \frac{b'(r)}{r^2} \right) \right) H(r) = \tilde{E}_{1,l}^2 H(r) \\ \left(-\Delta_l + 2 \left(\frac{b'(r)}{2r^2} + \frac{b(r)}{2r^3} - \frac{b'(r)}{r^2} \right) \right) K(r) = \tilde{E}_{2,l}^2 K(r) \end{cases}, \quad (66)$$

where Δ_l is

$$\Delta_l = \left(1 - \frac{b(r)}{r} \right) \frac{d^2}{dr^2} + \left(\frac{4r - b'(r)r - 3b(r)}{2r^2} \right) \frac{d}{dr} - \frac{l(l+1)}{r^2} - \frac{6}{r^2} \left(1 - \frac{b(r)}{r} \right). \quad (67)$$

Defining reduced fields

$$H(r) = \frac{f_1(r)}{r}; \quad K(r) = \frac{f_2(r)}{r}, \quad (68)$$

and passing to the proper geodesic distance from the *throat* of the bridge defined by

$$dx = \pm \frac{dr}{\sqrt{1 - \frac{b(r)}{r}}}, \quad (69)$$

the system (66) becomes ($r \equiv r(x)$)

$$\begin{cases} \left[-\frac{d^2}{dx^2} + V_1(r) \right] f_1(x) = \tilde{E}_{1,l}^2 f_1(x) \\ \left[-\frac{d^2}{dx^2} + V_2(r) \right] f_2(x) = \tilde{E}_{2,l}^2 f_2(x) \end{cases} \quad (70)$$

where we have defined $r \equiv r(x)$ and

$$\begin{cases} V_1(r) = \frac{l(l+1)}{r^2} + U_1(r) \\ V_2(r) = \frac{l(l+1)}{r^2} + U_2(r) \end{cases}, \quad (71)$$

with

$$\begin{cases} U_1(r) = c_1(r) + \left(\frac{1}{\omega} - 3 \right) c_2(r), \\ U_2(r) = c_1(r) + 3 \left(\frac{1}{\omega} + 1 \right) c_2(r), \\ c_1(r) = \frac{6}{r^2} \left(1 - \left(\frac{r_*}{r} \right)^{1+\frac{1}{\omega}} \right) \quad c_2(r) = \frac{1}{2r^2} \left(\frac{r_*}{r} \right)^{1+\frac{1}{\omega}} \end{cases}. \quad (72)$$

In order to use the WKB approximation, we define two r -dependent radial wave numbers $k_1(x, l, \tilde{E}_{1,nl})$ and $k_2(x, l, \tilde{E}_{2,nl})$

$$\begin{cases} k_1^2(x, l, \tilde{E}_{1,nl}) = \tilde{E}_{1,nl}^2 - \frac{l(l+1)}{r^2} - U_1(r) \\ k_2^2(x, l, \tilde{E}_{2,nl}) = \tilde{E}_{2,nl}^2 - \frac{l(l+1)}{r^2} - U_2(r) \end{cases}. \quad (73)$$

The number of modes with frequency less than $\tilde{E}_i$, $i = 1, 2$, is given approximately by

$$\tilde{g}(\tilde{E}_i) = \int \nu_i(l, \tilde{E}_i) (2l+1) dl, \quad (74)$$

where $\nu_i(l, \omega_i)$, $i = 1, 2$ is the number of nodes in the mode with (l, ω_i) , such that

$$\nu_i(l, \omega_i) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} dx \sqrt{k_i^2(x, l, \omega_i)}. \quad (75)$$

Here it is understood that the integration with respect to x and l is taken over those values which satisfy $k_i^2(x, l, \tilde{E}_i) \geq 0$, $i = 1, 2$. Thus the total one loop energy for TT tensors is given by (recall that $r \equiv r(x)$)

$$\begin{aligned} E^{TT} &= \frac{1}{4} \sum_{i=1}^2 \int_0^{+\infty} \tilde{E}_i \frac{d\tilde{g}(\tilde{E}_i)}{d\tilde{E}_i} d\tilde{E}_i = \sum_{i=1}^2 \int_{-\infty}^{+\infty} dx r^2 \left[\frac{1}{4\pi} \int_{\sqrt{U_i(r)}}^{+\infty} \tilde{E}_i^2 \sqrt{\tilde{E}_i^2 - U_i(r)} d\tilde{E}_i \right] \\ &= \int_{r_t}^{+\infty} dr r^2 [\rho_1 + \rho_2], \end{aligned} \quad (76)$$

where

$$\begin{cases} \rho_1 = \frac{1}{4\pi} \int_{\sqrt{U_1(x)}}^{+\infty} \tilde{E}_1^2 \sqrt{\tilde{E}_1^2 - U_1(r)} d\tilde{E}_1 \\ \rho_2 = \frac{1}{4\pi} \int_{\sqrt{U_2(x)}}^{+\infty} \tilde{E}_2^2 \sqrt{\tilde{E}_2^2 - U_2(r)} d\tilde{E}_2 \end{cases}. \quad (77)$$

VI. ONE LOOP ENERGY REGULARIZATION AND RENORMALIZATION

In this section, we proceed to evaluate the one loop energy. The method is equivalent to the scattering phase shift method and to the same method used to compute the entropy in the brick wall model. We use the zeta function regularization method to compute the energy densities ρ_1 and ρ_2 . Note that this procedure is completely equivalent to the subtraction procedure of the Casimir energy computation where zero point energy (ZPE) in different backgrounds with the same asymptotic properties is involved. To this purpose, we introduce the additional mass parameter μ in order to restore the correct dimension for the regularized quantities. Such an arbitrary mass scale emerges unavoidably in any regularization schemes. Then we have

$$\rho_i(\varepsilon) = \frac{1}{4\pi} \mu^{2\varepsilon} \int_{\sqrt{U_i(r)}}^{+\infty} d\tilde{E}_i \frac{\tilde{E}_i^2}{(\tilde{E}_i^2 - U_i(r))^{\varepsilon - \frac{1}{2}}} \quad (78)$$

If one of the functions $U_i(r)$ is negative, then the integration has to be meant in the range where $\tilde{E}_i^2 + U_i(r) \geq 0$. In both cases the result of the integration is⁵

$$= -\frac{U_i^2(r)}{64\pi^2} \left[\frac{1}{\varepsilon} + \ln \left(\frac{\mu^2}{U_i(r)} \right) + 2 \ln 2 - \frac{1}{2} \right], \quad (79)$$

where the absolute value has been inserted to take account of the possible change of sign. Then the total regularized one loop energy is

$$E^{TT}(r_t, \varepsilon; \mu) = 4\pi \left\{ 2 \int_{r_t}^{+\infty} dr \frac{r^2}{\sqrt{1 - \frac{b(r)}{r}}} [(\rho_1(\varepsilon) + \rho_2(\varepsilon))] \right\}, \quad (80)$$

where the factor 4π comes from the angular integration, while the factor 2 in front of the integral appears because we have come back to the original radial coordinate r : this means that we have to double the computation because of the upper and lower universe. To further proceed, it is useful to define the following coefficients:

⁵ Details of the calculation can be found in the Appendix D.

1.

$$a := 2r_t \int_{r_t}^{+\infty} dr \frac{r^2}{\sqrt{1 - \frac{b(r)}{r}}} [U_1^2(r) + U_2^2(r)], \quad (81)$$

and

2.

$$\tilde{b} := -2r_t \int_{r_t}^{+\infty} dr \frac{r^2}{\sqrt{1 - \frac{b(r)}{r}}} [U_1^2(r) \ln |U_1(r)| + U_2^2(r) \ln (U_2(r))]. \quad (82)$$

$$c := 2r_t \int_{r_t}^{+\infty} dr \frac{r^2}{\sqrt{1 - \frac{b(r)}{r}}} [U_1^2(r) + U_2^2(r)] \ln(\mu^2) = 2a \ln \mu. \quad (83)$$

Rescale the radial coordinate r so that $\rho = r/r_t$, then

$$\frac{b(r)}{r} = \left(\frac{r_t}{r}\right)^{1+\frac{1}{\omega}} = \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}; \quad U_1(r) = U_1(\rho)/r_t^2; \quad U_2(r) = U_2(\rho)/r_t^2. \quad (84)$$

Thus the coefficient a simply becomes

$$a := 2 \int_1^{+\infty} d\rho \frac{\rho^2}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} [U_1^2(\rho) + U_2^2(\rho)], \quad (85)$$

while the coefficient $\tilde{b}$ changes into

$$\begin{aligned} \tilde{b} &:= -2 \int_1^{+\infty} d\rho \frac{\rho^2}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} [U_1^2(\rho) \ln |U_1(\rho)| + U_2^2(\rho) \ln (U_2(\rho))] + 2 \int_1^{+\infty} d\rho \frac{\rho^2}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} [U_1^2(\rho) + U_2^2(\rho)] (2 \ln r_t) \\ &= -2 \int_1^{+\infty} d\rho \frac{\rho^2}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} [U_1^2(\rho) \ln |U_1(\rho)| + U_2^2(\rho) \ln (U_2(\rho))] + 2a \ln r_t = b + 2a \ln r_t. \end{aligned}$$

The result of the integration over the r coordinate leads to the following expression

$$E^{TT}(r_t, \varepsilon; \mu) = \frac{1}{16\pi} \left[-\frac{a}{\varepsilon r_t} - \frac{b}{r_t} - \frac{2a}{r_t} \ln \left(\frac{\sqrt{8} r_t \mu}{\sqrt[4]{e}} \right) \right] \quad (86)$$

and the self consistent equation (55) can be written in the form

$$A(\omega) \frac{r_t}{G} = \frac{1}{16\pi} \left[\frac{a}{\varepsilon r_t} + \frac{b}{r_t} + \frac{2a}{r_t} \ln \left(\frac{\sqrt{8} r_t \mu}{\sqrt[4]{e}} \right) \right]. \quad (87)$$

Actually, also the coefficients a and b depend on ω , but at this level is not relevant. In order to deal with finite quantities, we renormalize the divergent energy by absorbing the singularity in the classical quantity. In particular, we re-define the bare classical constant G

$$\frac{1}{G} \rightarrow \frac{1}{G_0} + \frac{a}{\varepsilon A(\omega) 16\pi r_t^2}. \quad (88)$$

Therefore, the remaining finite value for the effective equation (87) reads

$$A(\omega) \frac{r_t}{G_0} = \frac{1}{16\pi} \left[\frac{b}{r_t} + \frac{2a}{r_t} \ln \left(\frac{\sqrt{8} r_t \mu}{\sqrt[4]{e}} \right) \right]. \quad (89)$$

This quantity depends on the arbitrary mass scale μ . It is appropriate to use the renormalization group equation to eliminate such a dependence. To this aim, we impose that[21]

$$\mu \frac{d}{d\mu} \left(\frac{A(\omega) r_t}{G_0(\mu)} \right) = \mu \frac{d}{d\mu} \left\{ \frac{1}{16\pi} \left[\frac{b}{r_t} + \frac{2a}{r_t} \ln \left(\frac{\sqrt{8} r_t \mu}{\sqrt[4]{e}} \right) \right] \right\}, \quad (90)$$

namely

$$A(\omega) r_t \mu \frac{\partial G_0^{-1}(\mu)}{\partial \mu} - \frac{a}{8\pi r_t} = 0. \quad (91)$$

Solving it we find that the renormalized constant G_0 should be treated as a running one in the sense that it varies provided that the scale μ is changing

$$\frac{1}{G_0(\mu)} = \frac{1}{G_0(\mu_0)} + K \ln \left(\frac{\mu}{\mu_0} \right) \quad \text{or} \quad G_0(\mu) = \frac{G_0(\mu_0)}{1 + G_0(\mu_0) K \ln \left(\frac{\mu}{\mu_0} \right)}, \quad K = \frac{a}{A(\omega) 8\pi r_t^2}; \quad (92)$$

where μ_0 is the normalization point. We substitute Eq.(92) into Eq.(89) to find

$$\frac{A(\omega)}{G_0(\mu_0)} = \frac{1}{16\pi} \left[\frac{b}{r_t^2} + \frac{2a}{r_t^2} \ln \left(\frac{\sqrt{8} r_t \mu_0}{\sqrt[4]{e}} \right) \right], \quad (93)$$

where we have divided by r_t . In order to have only one solution⁶, we find the extremum of the r.h.s. of Eq.(93) and we get

$$\frac{a-b}{2a} = \ln \left(\frac{\sqrt{8} \bar{r}_t \mu_0}{\sqrt[4]{e}} \right) \quad \Rightarrow \quad \bar{r}_t = \frac{\sqrt[4]{e}}{\sqrt{8} \mu_0} \exp \left(\frac{a-b}{2a} \right) \quad (94)$$

and

$$\frac{1}{G_0(\mu_0)} = \frac{a \mu_0^2}{2\pi A(\omega) \sqrt{e}} \exp \left(-\frac{a-b}{a} \right). \quad (95)$$

With the help of Eqs.(94) and (95), Eq.(92) becomes

$$\begin{aligned} \frac{1}{G_0(\mu)} &= \frac{1}{G_0(\mu_0)} + \frac{a}{A(\omega) 8\pi \bar{r}_t^2} \ln \left(\frac{\mu}{\mu_0} \right) = \frac{a \mu_0^2}{A(\omega) 2\pi \sqrt{e}} \exp \left(-\frac{a-b}{a} \right) \left[1 + 2 \ln \left(\frac{\mu}{\mu_0} \right) \right] \\ &= \frac{1}{G_0(\mu_0)} \left[1 + 2 \ln \left(\frac{\mu}{\mu_0} \right) \right]. \end{aligned} \quad (96)$$

It is straightforward to see that we have a constraint on μ/μ_0 . Indeed we have to choose

$$\mu > \mu_0 \exp \left(-\frac{1}{2} \right) = .6065306597 \mu_0, \quad (97)$$

otherwise $G_0(\mu)$ becomes negative[16]. We have now two possibilities: 1) we identify $G_0(\mu_0)$ with the squared Planck length, then the wormhole radius becomes

$$\bar{r}_t = \sqrt{\frac{a(\omega)}{16\pi A(\omega)}} l_p, \quad (98)$$

⁶ Note that in the paper of Khusnutdinov and Sushkov[11], to find only one solution, the minimum of the ground state of the quantized scalar field has been set equal to the classical energy. In our case, we have no external fields on a given background. This means that it is not possible to find a minimum of the one loop gravitons, in analogy with Ref.[11]. Moreover the renormalization procedure in Ref.[11] is completely independent by the classical term, while in our case it is not. Indeed, thanks to the self-consistent equation (55), we can renormalize the divergent term.

where we have reestablished the ω dependence of the coefficient a . It is useful to write the expression for $\omega \rightarrow +\infty$ and for $\omega \rightarrow 0$. We get

$$\left\{ \begin{array}{ll} \bar{r}_t \simeq \left[\frac{\sqrt{105}}{5\sqrt{\pi}} \left(1 + \frac{1}{\omega} \left(\frac{449}{420} - 2 \ln(2) \right) + O(\omega^{-2}) \right) \right] l_p & \omega \rightarrow +\infty \\ \bar{r}_t \simeq \left[\frac{\sqrt{30}}{12\sqrt{\pi}\sqrt{\omega}} + O(\omega^{1/2}) \right] l_p & \omega \rightarrow 0 \end{array} \right. . \quad (99)$$

The following plot shows the behavior of $\bar{r}_t$ as a function of ω . It is visible the presence of a minimum for $\bar{\omega} = 3.35204$,





FIG. 1: Plot of the wormhole throat $\bar{r}_t$ as a function of ω in the positive range.

where $\bar{r}_t(\bar{\omega}) = 1.11891$. As we can see, from the expression (99) and from the Fig.1, the radius is divergent when $\omega \rightarrow 0$. At this stage, we cannot establish if this is a physical result or a failure of the scheme. When $\omega \rightarrow \pm\infty$, $\bar{r}_t$ approaches the value $1.15624l_p$, while for $\omega = 1$, we obtained $\bar{r}_t = 1.15882l_p$. It is interesting to note that when $\omega \rightarrow +\infty$, the shape function $b(r)$ in Eq.(23) approaches the Schwarzschild value, when we identify $\bar{r}_t$ with $2MG$. In this sense, it seems that also the Schwarzschild wormhole is traversable.

2) We identify μ_0 with the Planck scale and we get from Eq.(94) the following plot Note the absence of a minimum.



Differently from case of Eq.(98), there is no access to the phantom range. In our case, we suppose that the graviton quantum fluctuations play the role of the exotic matter and even if we fix the renormalization point at the Planck scale as in Ref.[11], we find a radius $\bar{r}_t > r_w$.

VII. SUMMARY AND CONCLUSIONS

In this paper, we have generalized the analysis of self-sustaining wormholes[7] by looking how the equation of state (2) can affect the traversability, when the sign of the parameter ω is positive. The paper has been motivated by the work of Lobo[4], Kuhfittig[5] and Sushkov[6], where the authors search for classical traversable wormholes supported by phantom energy. Since the phantom energy must satisfy the equation of state, but in the range $\omega < -1$, we have investigated the possibility of studying the whole range $-\infty < \omega < +\infty$. Unfortunately, evaluating the classical term we have discovered that such a term is well defined in the range $-1 < \omega < +\infty$. The interval $-1 < \omega < 0$ should be interesting for the existence of a “dark” energy support. Once again, the “dark” energy domain lies outside the asymptotically flatness property. So, unless one is interested in wormholes that are not asymptotically flat, i.e. asymptotically de Sitter or asymptotically Anti-de Sitter, we have to reject also this possibility. Therefore, the final stage of computation has been restricted only to positive values of the parameter ω . In this context, it is interesting to note that also the Schwarzschild wormhole is traversable, even if in the limiting procedure of $\omega \rightarrow +\infty$. Despite of this, the obtained “traversability” has to be regarded as in “principle” rather than in “practice” because the wormhole radius has a Planckian size. We do not know, at this stage of the calculation, if a different approach for a self sustained wormhole can give better results. On the other hand, the positive ω sector seems to corroborate the Casimir process of the quantum fluctuations supporting the opening of the wormhole. Even in this region, we do not know what happens approaching directly the point $\omega = 0$, because it seems that this approach is ill defined. Nevertheless, in this paper we have studied the behavior of the energy. Work in progress seems to show that dealing with energy density one can get more general results even in the “phantom” sector[8].

Acknowledgments

The author would like to thank S. Capozziello, F. Lobo and the Referee for useful comments and suggestions. In particular, the author would like to thank the Referee, for having brought to his attention the paper of Ref.[6].

APPENDIX A: COMPUTATION OF THE CLASSICAL TERM

Here, we give details leading to Eqs.(53, 54). We begin with the definition of the hamiltonian which in the static case simplifies into

$$\int_{\Sigma} d^3x \mathcal{H}^{(0)} = -\frac{1}{16\pi G} \int_{\Sigma} d^3x \sqrt{g} R^{(3)}. \quad (\text{A1})$$

With the help of Eqs(23, 24) , we get

$$\frac{1}{G\omega} \int_{r_t}^{+\infty} dr \left(\frac{r_t}{r} \right)^{\frac{x}{2}} \frac{1}{\sqrt{\left(\frac{r}{r_t} \right)^x - 1}} \quad x := 1 + \frac{1}{\omega}. \quad (\text{A2})$$

In the previous integral there is an extra factor “2” coming from the counting of the universes. We change variable to obtain

$$\frac{2r_t}{G\omega x} \int_0^{+\infty} \frac{dt}{\cosh^{2-2/x}(t)} = \frac{r_t}{G(1+\omega)} B\left(\frac{1}{2}, \frac{1}{1+\omega}\right) = \frac{r_t}{G} A(\omega) \quad (\text{A3})$$

with

$$A(\omega) = \frac{1}{(1+\omega)} B\left(\frac{1}{2}, \frac{1}{1+\omega}\right) = \frac{\sqrt{\pi}}{(1+\omega)} \frac{\Gamma\left(\frac{1}{1+\omega}\right)}{\Gamma\left(\frac{3+\omega}{2+2\omega}\right)}. \quad (\text{A4})$$

In Eq.(A3), we have used the following formula

$$\int_0^{+\infty} dt \frac{\sinh^{\mu} t}{\cosh^{\nu} t} = \frac{1}{2} B\left(\frac{\mu+1}{2}, \frac{\nu-\mu}{2}\right) \quad \left\{ \begin{array}{l} \text{Re } \mu > -1 \\ \text{Re } (\mu - \nu) < 0 \end{array} \right. . \quad (\text{A5})$$

APPENDIX B: EINSTEIN EQUATIONS AND THE HAMILTONIAN

Let us consider the Einstein equations

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = \kappa 8\pi G T_{\mu\nu}. \quad (\text{B1})$$

$R_{\mu\nu}$ is the Ricci tensor and R is the scalar curvature. If u^{μ} is a time-like unit vector such that $g_{\mu\nu} u^{\mu} u^{\nu} = -1$, then the Einstein tensor $G_{\mu\nu}$ becomes

$$G_{\mu\nu} u^{\mu} u^{\nu} = R_{\mu\nu} u^{\mu} u^{\nu} - \frac{1}{2} g_{\mu\nu} u^{\mu} u^{\nu} R = R_{\mu\nu} u^{\mu} u^{\nu} + \frac{1}{2} R. \quad (\text{B2})$$

By means of the Gauss-Codazzi equations[22],

$$R = R^{(3)} \pm 2R_{\mu\nu} u^{\mu} u^{\nu} \mp K^2 \pm K_{\mu\nu} K^{\mu\nu}, \quad (\text{B3})$$

where $K_{\mu\nu}$ is the extrinsic curvature and $R^{(3)}$ is the three dimensional scalar curvature. For a time-like vector, we take the lower sign and Eq.(B2) becomes

$$G_{\mu\nu} u^{\mu} u^{\nu} = \frac{1}{2} \left(R^{(3)} + K^2 - K_{\mu\nu} K^{\mu\nu} \right). \quad (\text{B4})$$

If the conjugate momentum is defined by

$$\pi^{\mu\nu} = \frac{\sqrt{{}^{(3)}g}}{2\kappa} (K g^{\mu\nu} - K^{\mu\nu}), \quad (\text{B5})$$

then

$$K^2 - K_{\mu\nu}K^{\mu\nu} = \left(\frac{2\kappa}{\sqrt{(3)g}} \right)^2 \left(\frac{\pi^2}{2} - \pi^{\mu\nu}\pi_{\mu\nu} \right) \quad (\text{B6})$$

and

$$\frac{\sqrt{(3)g}}{2\kappa} G_{\mu\nu} u^\mu u^\nu = \frac{\sqrt{(3)g}}{2\kappa} R^{(3)} + \frac{2\kappa}{\sqrt{(3)g}} \left(\frac{\pi^2}{2} - \pi^{\mu\nu}\pi_{\mu\nu} \right) = -\mathcal{H}^{(0)}, \quad (\text{B7})$$

namely Eq.(10) with the reversed sign.

APPENDIX C: ENERGY COEFFICIENTS

In this appendix, we report details on computation of the coefficients of the one loop graviton energy. We begin with the coefficient a defined by

$$a := r_t \int_{r_t}^{+\infty} dr \frac{r^2}{\sqrt{1 - \frac{b(r)}{r}}} [U_1^2(r) + U_2^2(r)], \quad (\text{C1})$$

which, with the help of Eqs. (72), becomes

$$a = r_t \int_{r_t}^{+\infty} dr \frac{r^2}{\sqrt{1 - \left(\frac{r_t}{r}\right)^{1+\frac{1}{\omega}}}} \left[2c_1^2(r) + 2 \left(9 + \frac{6}{\omega} + \frac{5}{\omega^2} \right) c_2^2(r) + \frac{8}{\omega} c_1(r) c_2(r) \right], \quad (\text{C2})$$

where

$$c_1(r) = \frac{6}{r^2} \left(1 - \left(\frac{r_t}{r} \right)^{1+\frac{1}{\omega}} \right) \quad c_2(r) = \frac{1}{2r^2} \left(\frac{r_t}{r} \right)^{1+\frac{1}{\omega}}. \quad (\text{C3})$$

Defining the dimensionless variable $\rho = \frac{r}{r_t}$, we can write

$$\begin{aligned} a_1 &= r_t \int_{r_t}^{+\infty} dr \frac{r^2}{\sqrt{1 - \left(\frac{r_t}{r}\right)^{1+\frac{1}{\omega}}}} [2c_1^2(r)] = 72 \int_1^{+\infty} d\rho \left(\rho^{(1+\frac{1}{\omega})} - 1 \right)^{\frac{3}{2}} \rho^{-2-\frac{3}{2}(1+\frac{1}{\omega})} \\ &= 72 \frac{\omega}{1+\omega} B \left(\frac{5}{2}, \frac{\omega}{1+\omega} \right). \end{aligned} \quad (\text{C4})$$

$$\begin{aligned} a_2 &= r_t \int_{r_t}^{+\infty} dr \frac{r^2}{\sqrt{1 - \left(\frac{r_t}{r}\right)^{1+\frac{1}{\omega}}}} \left[2 \left(9 + \frac{6}{\omega} + \frac{5}{\omega^2} \right) c_2^2(r) \right] = \frac{1}{2} \left(9 + \frac{6}{\omega} + \frac{5}{\omega^2} \right) \int_1^{+\infty} d\rho \left(\rho^{(1+\frac{1}{\omega})} - 1 \right)^{-\frac{1}{2}} \rho^{-2-\frac{3}{2}(1+\frac{1}{\omega})} \\ &= \frac{1}{2} \left(9 + \frac{6}{\omega} + \frac{5}{\omega^2} \right) \frac{\omega}{1+\omega} B \left(\frac{1}{2}, \frac{3\omega+2}{1+\omega} \right) \end{aligned} \quad (\text{C5})$$

and

$$\begin{aligned} a_3 &= r_t \int_{r_t}^{+\infty} dr \frac{r^2}{\sqrt{1 - \left(\frac{r_t}{r}\right)^{1+\frac{1}{\omega}}}} \left[\frac{8}{\omega} c_1(r) c_2(r) \right] = \frac{24}{\omega} \int_1^{+\infty} d\rho \left(\rho^{(1+\frac{1}{\omega})} - 1 \right)^{\frac{1}{2}} \rho^{-2-\frac{3}{2}(1+\frac{1}{\omega})} \\ &= \frac{24}{1+\omega} B \left(\frac{3}{2}, \frac{2\omega+1}{1+\omega} \right), \end{aligned} \quad (\text{C6})$$

so that

$$a \equiv a(\omega) = \frac{\omega}{1+\omega} \left[72B\left(\frac{5}{2}, \frac{\omega}{1+\omega}\right) + \frac{1}{2} \left(9 + \frac{6}{\omega} + \frac{5}{\omega^2}\right) B\left(\frac{1}{2}, \frac{3\omega+2}{1+\omega}\right) + \frac{24}{\omega} B\left(\frac{3}{2}, \frac{2\omega+1}{1+\omega}\right) \right], \quad (\text{C7})$$

where $B(x, y)$ is the beta function.. The same procedure applies to the coefficient b .

$$b := -2 \int_1^{+\infty} d\rho \frac{\rho^2}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} \left[U_1^2(\rho) \ln |U_1(\rho)| + U_2^2(\rho) \ln (U_2(\rho)) \right]. \quad (\text{C8})$$

We can separate the logarithmic functions

$$\begin{cases} \ln |U_1(\rho)| = \underbrace{\ln \left| 12\omega \left(\rho^{(1+\frac{1}{\omega})} - 1 \right) + (1-3\omega) \right|}_{b_{1a}} - \underbrace{\ln(2\omega)}_{b_{1b}} - \underbrace{\left(3 + \frac{1}{\omega} \right) \ln \rho}_{b_{1c}} \\ \ln |U_2(\rho)| = \underbrace{\ln \left| 12\omega \left(\rho^{(1+\frac{1}{\omega})} - 1 \right) + 3(1+\omega) \right|}_{b_{2a}} - \underbrace{\ln(2\omega)}_{b_{2b}} - \underbrace{\left(3 + \frac{1}{\omega} \right) \ln \rho}_{b_{2c}} \end{cases} \quad (\text{C9})$$

and compute separately the different terms. $b_{1b} + b_{2b}$ simply becomes

$$b_{1b} + b_{2b} = 2 \ln(2\omega) \int_1^{+\infty} d\rho \frac{\rho^2 [U_1^2(\rho) + U_2^2(\rho)]}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} = \ln(2\omega) a(\omega). \quad (\text{C10})$$

The coefficient $b_{1c} + b_{2c}$ is

$$b_{1c} + b_{2c} = \left(3 + \frac{1}{\omega} \right) \int_1^{+\infty} d\rho \frac{\rho^2 [U_1^2(\rho) + U_2^2(\rho)]}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} \ln \rho, \quad (\text{C11})$$

namely

$$\left(3 + \frac{1}{\omega} \right) \int_1^{+\infty} d\rho \frac{\rho^2}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} \left[2c_1^2(\rho) + 2 \left(9 + \frac{6}{\omega} + \frac{5}{\omega^2} \right) c_2^2(\rho) + \frac{8}{\omega} c_1(\rho) c_2(\rho) \right] \ln \rho \quad (\text{C12})$$

The first term, $b_{f,1}$ will be summed with the coefficient a , while the second term can be written as $b_{f,b1} + b_{f,b2} + b_{f,b3}$ with

$$\begin{aligned} 2 \left(3 + \frac{1}{\omega} \right) \int_1^{+\infty} d\rho \frac{\rho^2 c_1^2(\rho)}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} \ln \rho &= 72 \left(3 + \frac{1}{\omega} \right) \left(\frac{\omega}{1+\omega} \right)^2 \int_1^{+\infty} d\rho \left(\rho^{(1+\frac{1}{\omega})} - 1 \right)^{\frac{3}{2}} \rho^{-2-\frac{3}{2}(1+\frac{1}{\omega})} \ln \rho \\ &= 72 \left(3 + \frac{1}{\omega} \right) \left(\frac{\omega}{1+\omega} \right)^2 \int_0^{+\infty} ds s^{\frac{3}{2}} (1+s)^{-\frac{7\omega+5}{2(1+\omega)}} \ln(1+s) \\ &= 72 \left(3 + \frac{1}{\omega} \right) \left(\frac{\omega}{1+\omega} \right)^2 B\left(\frac{5}{2}, \frac{\omega}{1+\omega}\right) \left[\Psi\left(\frac{7\omega+5}{2(1+\omega)}\right) - \Psi\left(\frac{\omega}{1+\omega}\right) \right]. \quad (\text{C13}) \\ &\quad \left(3 + \frac{1}{\omega} \right) 2 \left(9 + \frac{6}{\omega} + \frac{5}{\omega^2} \right) \int_1^{+\infty} d\rho \frac{\rho^2 c_2^2(\rho)}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} \ln \rho \end{aligned}$$

$$\begin{aligned}
&= \left(3 + \frac{1}{\omega}\right) \frac{1}{2} \left(9 + \frac{6}{\omega} + \frac{5}{\omega^2}\right) \int_1^{+\infty} d\rho \left(\rho^{(1+\frac{1}{\omega})} - 1\right)^{-\frac{1}{2}} \rho^{-2-\frac{3}{2}(1+\frac{1}{\omega})} \ln \rho \\
&= \left(3 + \frac{1}{\omega}\right) \frac{1}{2} \left(9 + \frac{6}{\omega} + \frac{5}{\omega^2}\right) \left(\frac{\omega}{1+\omega}\right)^2 \int_0^{+\infty} ds s^{-\frac{1}{2}} (1+s)^{-\frac{7\omega+5}{2(1+\omega)}} \ln(1+s) \\
&= \left(3 + \frac{1}{\omega}\right) \frac{1}{2} \left(9 + \frac{6}{\omega} + \frac{5}{\omega^2}\right) \left(\frac{\omega}{1+\omega}\right)^2 B\left(\frac{1}{2}, \frac{3\omega+2}{1+\omega}\right) \left[\Psi\left(\frac{7\omega+5}{2(1+\omega)}\right) - \Psi\left(\frac{3\omega+2}{1+\omega}\right)\right] \quad (C14)
\end{aligned}$$

and

$$\begin{aligned}
\left(3 + \frac{1}{\omega}\right) \frac{8}{\omega} \int_1^{+\infty} d\rho \frac{\rho^2 c_1(\rho) c_2(\rho)}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} \ln \rho &= \left(3 + \frac{1}{\omega}\right) \frac{24}{\omega} \int_1^{+\infty} d\rho \left(\rho^{(1+\frac{1}{\omega})} - 1\right)^{\frac{1}{2}} \rho^{-2-\frac{3}{2}(1+\frac{1}{\omega})} \ln \rho \\
&= \left(3 + \frac{1}{\omega}\right) \frac{24}{\omega} \left(\frac{\omega}{1+\omega}\right)^2 \int_0^{+\infty} ds s^{\frac{1}{2}} (1+s)^{-\frac{7\omega+5}{2(1+\omega)}} \ln(1+s) \\
&= \left(3 + \frac{1}{\omega}\right) \frac{24}{\omega} \left(\frac{\omega}{1+\omega}\right)^2 B\left(\frac{3}{2}, \frac{2\omega+1}{1+\omega}\right) \left[\Psi\left(\frac{7\omega+5}{2(1+\omega)}\right) - \Psi\left(\frac{2\omega+1}{1+\omega}\right)\right], \quad (C15)
\end{aligned}$$

where we have used the following relation

$$\int_0^{+\infty} dx x^{\mu-1} \frac{\ln(\gamma+x)}{(\gamma+x)^\nu} = \gamma^{\mu-\nu} B(\mu, \nu-\mu) [\Psi(\nu) - \Psi(\nu-\mu) + \ln \gamma], \quad (C16)$$

where Ψ is the digamma function. Globally, the coefficient $b_{1c} + b_{2c}$ becomes

$$\begin{aligned}
&= \left(3 + \frac{1}{\omega}\right) \left(\frac{\omega}{1+\omega}\right)^2 \left\{ 72B\left(\frac{5}{2}, \frac{\omega}{1+\omega}\right) \left[\Psi\left(\frac{7\omega+5}{2(1+\omega)}\right) - \Psi\left(\frac{\omega}{1+\omega}\right)\right] \right. \\
&\quad \left. + \frac{24}{\omega} B\left(\frac{3}{2}, \frac{2\omega+1}{1+\omega}\right) \left[\Psi\left(\frac{7\omega+5}{2(1+\omega)}\right) - \Psi\left(\frac{2\omega+1}{1+\omega}\right)\right] \right\}. \quad (C17)
\end{aligned}$$

The term $b_{1a} + b_{2b}$ is more complicated. The logarithmic term is of the form $\ln(\gamma_{1,2} + \delta x)$. It is useful to re-express it in terms of hypergeometric function, namely

$$\ln(\gamma_{1,2} + \delta x) = \delta \frac{x}{\gamma_{1,2}} {}_2F_1\left(1, 1; 2; -\delta \frac{x}{\gamma_{1,2}}\right) + \ln \gamma_{1,2}, \quad (C18)$$

with $\delta = 12\omega$ and

$$\begin{cases} \gamma_1 = 1 - 3\omega \\ \gamma_2 = 3(1 + \omega) \end{cases}. \quad (C19)$$

Expression

$$\ln \gamma_1 \int_1^{+\infty} d\rho \frac{\rho^2}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} \left[c_1^2(\rho) + \left(\frac{1}{\omega} - 3\right)^2 c_2^2(\rho) + 2\left(\frac{1}{\omega} - 3\right) c_1(\rho) c_2(\rho) \right] \quad (C20)$$

becomes

$$\ln \gamma_1 \frac{\omega}{1+\omega} \left[36B\left(\frac{5}{2}, \frac{\omega}{1+\omega}\right) + \left(\frac{1}{\omega} - 3\right)^2 B\left(\frac{1}{2}, \frac{3\omega+2}{1+\omega}\right) + 6\left(\frac{1}{\omega} - 3\right) B\left(\frac{3}{2}, \frac{2\omega+1}{1+\omega}\right) \right] \quad (C21)$$

and expression

$$\ln \gamma_2 \int_1^{+\infty} d\rho \frac{\rho^2}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} \left[c_1^2(\rho) + \left(\frac{3}{\omega} + 3\right)^2 c_2^2(\rho) + 2 \left(\frac{3}{\omega} + 3\right) c_1(\rho) c_2(\rho) \right] \quad (C22)$$

becomes

$$\ln \gamma_2 \frac{\omega}{1+\omega} \left[36B\left(\frac{5}{2}, \frac{\omega}{1+\omega}\right) + \left(\frac{3}{\omega} + 3\right)^2 B\left(\frac{1}{2}, \frac{3\omega+2}{1+\omega}\right) + 6 \left(\frac{3}{\omega} + 3\right) B\left(\frac{3}{2}, \frac{2\omega+1}{1+\omega}\right) \right]. \quad (C23)$$

Concerning the hypergeometric part, after having defined the variable $s = \rho^{(1+\frac{1}{\omega})} - 1$, we use the following formula

$$\int_0^{+\infty} dx \frac{x^{\alpha-1}}{(x+z)^\rho} {}_2F_1(a, b; c; -\omega x) dx = z^{\alpha-\rho} B(\alpha, \rho-\alpha) {}_3F_2(a, b, \alpha; c, \alpha-\rho+1; \omega z) \\ + \omega^{\rho-\alpha} \frac{\Gamma(c) \Gamma(a-\alpha+\rho) \Gamma(b-\alpha+\rho) \Gamma(\alpha-\rho)}{\Gamma(a) \Gamma(b) \Gamma(c-\alpha+\rho)} {}_3F_2(a-\alpha+\rho, b-\alpha+\rho, \rho; c-\alpha+\rho, \rho-\alpha+1; \omega z), \quad (C24)$$

which, in our specific case, becomes

$$\frac{\delta}{\gamma_{1,2}} \int_0^{+\infty} ds \frac{s^{\alpha-1}}{(s+1)^\rho} s {}_2F_1\left(1, 1; 2; -\delta \frac{s}{\gamma_{1,2}}\right) ds = B(\alpha, \rho-\alpha) {}_3F_2\left(1, 1, \alpha; 2, \alpha-\rho+1; \frac{\delta}{\gamma_{1,2}}\right) \\ + \left(\frac{\delta}{\gamma_{1,2}}\right)^{\rho-\alpha} \frac{\Gamma(1-\alpha+\rho) \Gamma(1-\alpha+\rho) \Gamma(\alpha-\rho)}{\Gamma(2-\alpha+\rho)} {}_3F_2\left(1-\alpha+\rho, 1-\alpha+\rho, \rho; 2-\alpha+\rho, 1-\alpha+\rho; \frac{\delta}{\gamma_{1,2}}\right). \quad (C25)$$

Therefore for γ_1 we get

$$\frac{\delta}{\gamma_1} \int_1^{+\infty} d\rho \frac{{}_2F_1\left(1, 1; 2; -\frac{\delta}{\gamma_1} \left(\rho^{(1+\frac{1}{\omega})} - 1\right)\right) \rho^2}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} \left(\rho^{(1+\frac{1}{\omega})} - 1\right) \left[c_1^2(\rho) + \left(\frac{1}{\omega} - 3\right)^2 c_2^2(\rho) + 2 \left(\frac{1}{\omega} - 3\right) c_1(\rho) c_2(\rho) \right]. \quad (C26)$$

Every piece leads to: a)

$$\frac{\delta}{\gamma_1} \int_1^{+\infty} d\rho \frac{\rho^2 c_1^2(\rho)}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} \left(\rho^{(1+\frac{1}{\omega})} - 1\right) {}_2F_1\left(1, 1; 2; -\frac{\delta}{\gamma_1} \left(\rho^{(1+\frac{1}{\omega})} - 1\right)\right) \\ = 36 \frac{\delta}{\gamma_1} \left(\frac{\omega}{1+\omega}\right) \int_0^{+\infty} ds s^{\frac{5}{2}} (1+s)^{-\frac{7\omega+5}{2(1+\omega)}} {}_2F_1\left(1, 1; 2; -\frac{\delta}{\gamma_1} s\right) \\ = 36 \frac{\delta}{\gamma_1} \left(\frac{\omega}{1+\omega}\right) \left[B\left(\frac{7}{2}, -\frac{2}{1+\omega}\right) {}_3F_2\left(1, 1, \frac{7}{2}; 2, \frac{\omega+3}{1+\omega}; \frac{\delta}{\gamma_1}\right) \right. \\ \left. + \left(\frac{\delta}{\gamma_1}\right)^{-\frac{2}{1+\omega}} \frac{\left[\Gamma\left(\frac{\omega-1}{1+\omega}\right)\right]^2 \Gamma\left(\frac{2}{1+\omega}\right)}{\Gamma\left(\frac{2\omega}{1+\omega}\right)} {}_3F_2\left(\frac{\omega-1}{1+\omega}, \frac{\omega-1}{1+\omega}, \frac{7\omega+5}{2(1+\omega)}; \frac{2\omega}{1+\omega}, \frac{\omega-1}{1+\omega}; \frac{\delta}{\gamma_1}\right) \right], \quad (C27)$$

b)

$$\frac{\delta}{\gamma_1} \left(\frac{1}{\omega} - 3\right)^2 \int_1^{+\infty} d\rho \frac{\rho^2 c_2^2(\rho)}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} \left(\rho^{(1+\frac{1}{\omega})} - 1\right) {}_2F_1\left(1, 1; 2; -\frac{\delta}{\gamma_1} \left(\rho^{(1+\frac{1}{\omega})} - 1\right)\right)$$

$$\begin{aligned}
&= \frac{\delta\gamma_1}{4\omega(1+\omega)} \int_0^{+\infty} ds s^{\frac{1}{2}} (1+s)^{-\frac{7\omega+5}{2(1+\omega)}} {}_2F_1\left(1, 1; 2; -\frac{\delta}{\gamma_1}s\right) = \frac{\delta\gamma_1}{4\omega(1+\omega)} \left[B\left(\frac{3}{2}, \frac{2\omega+1}{1+\omega}\right) {}_3F_2\left(1, 1, \frac{3}{2}; 2, -\frac{\omega}{1+\omega}; \frac{\delta}{\gamma_1}\right) \right. \\
&\quad \left. + \left(\frac{\delta}{\gamma_1}\right)^{-\frac{2\omega+1}{1+\omega}} \frac{\left[\Gamma\left(\frac{3\omega+2}{1+\omega}\right)\right]^2 \Gamma\left(-\frac{2\omega+1}{1+\omega}\right)}{\Gamma\left(\frac{4\omega+3}{1+\omega}\right)} {}_3F_2\left(\frac{3\omega+2}{1+\omega}, \frac{3\omega+2}{1+\omega}, \frac{7\omega+5}{2(1+\omega)}; \frac{4\omega+3}{1+\omega}, \frac{3\omega+2}{1+\omega}; \frac{\delta}{\gamma_1}\right) \right] \quad (C28)
\end{aligned}$$

and c)

$$\begin{aligned}
&\frac{\delta}{\gamma_1} \left(\frac{1}{\omega} - 3\right) \int_1^{+\infty} d\rho \frac{\rho^2 c_1(\rho) c_2(\rho)}{\sqrt{1 - \left(\frac{1}{\rho}\right)^{1+\frac{1}{\omega}}}} \left(\rho^{(1+\frac{1}{\omega})} - 1\right) {}_2F_1\left(1, 1; 2; -\frac{\delta}{\gamma_1} \left(\rho^{(1+\frac{1}{\omega})} - 1\right)\right) \\
&= \frac{6\delta}{1+\omega} \int_0^{+\infty} ds s^{\frac{3}{2}} (1+s)^{-\frac{7\omega+5}{2(1+\omega)}} {}_2F_1\left(1, 1; 2; -\frac{\delta}{\gamma_1}s\right) = \frac{6\delta}{1+\omega} \left[B\left(\frac{5}{2}, \frac{\omega}{1+\omega}\right) {}_3F_2\left(1, 1, \frac{5}{2}; 2, \frac{1}{1+\omega}; \frac{\delta}{\gamma_1}\right) \right. \\
&\quad \left. + \left(\frac{\delta}{\gamma_1}\right)^{\frac{\omega}{1+\omega}} \frac{\left[\Gamma\left(\frac{2\omega+1}{1+\omega}\right)\right]^2 \Gamma\left(-\frac{\omega}{1+\omega}\right)}{\Gamma\left(\frac{3\omega+2}{1+\omega}\right)} {}_3F_2\left(\frac{2\omega+1}{1+\omega}, \frac{2\omega+1}{1+\omega}, \frac{7\omega+5}{2(1+\omega)}; \frac{3\omega+2}{1+\omega}, \frac{2\omega+1}{1+\omega}; \frac{\delta}{\gamma_1}\right) \right]. \quad (C29)
\end{aligned}$$

The same set of expressions works for γ_2 too.

APPENDIX D: THE ZETA FUNCTION REGULARIZATION

In this appendix, we report details on computation leading to expression (78). We begin with the following integral

$$\rho(\varepsilon) = \begin{cases} I_+ = \mu^{2\varepsilon} \int_0^{+\infty} d\omega \frac{\omega^2}{(\omega^2 + U(x))^{\varepsilon - \frac{1}{2}}} \\ I_- = \mu^{2\varepsilon} \int_0^{+\infty} d\omega \frac{\omega^2}{(\omega^2 - U(x))^{\varepsilon - \frac{1}{2}}} \end{cases}, \quad (D1)$$

with $U(x) > 0$.

1. I_+ computation

If we define $t = \omega/\sqrt{U(x)}$, the integral I_+ in Eq.(D1) becomes

$$\begin{aligned}
\rho(\varepsilon) &= \mu^{2\varepsilon} U(x)^{2-\varepsilon} \int_0^{+\infty} dt \frac{t^2}{(t^2 + 1)^{\varepsilon - \frac{1}{2}}} = \frac{1}{2} \mu^{2\varepsilon} U(x)^{2-\varepsilon} B\left(\frac{3}{2}, \varepsilon - 2\right) \\
&= \frac{1}{2} \mu^{2\varepsilon} U(x)^{2-\varepsilon} \frac{\Gamma\left(\frac{3}{2}\right) \Gamma(\varepsilon - 2)}{\Gamma\left(\varepsilon - \frac{1}{2}\right)} = \frac{\sqrt{\pi}}{4} U(x)^2 \left(\frac{\mu^2}{U(x)}\right)^\varepsilon \frac{\Gamma(\varepsilon - 2)}{\Gamma\left(\varepsilon - \frac{1}{2}\right)}, \quad (D2)
\end{aligned}$$

where we have used the following identities involving the beta function

$$B(x, y) = 2 \int_0^{+\infty} dt \frac{t^{2x-1}}{(t^2 + 1)^{x+y}} \quad \text{Re } x > 0, \text{Re } y > 0 \quad (D3)$$

related to the gamma function by means of

$$B(x, y) = \frac{\Gamma(x) \Gamma(y)}{\Gamma(x+y)}. \quad (D4)$$

Taking into account the following relations for the Γ -function

$$\Gamma(\varepsilon - 2) = \frac{\Gamma(1 + \varepsilon)}{\varepsilon(\varepsilon - 1)(\varepsilon - 2)}, \quad \Gamma\left(\varepsilon - \frac{1}{2}\right) = \frac{\Gamma\left(\varepsilon + \frac{1}{2}\right)}{\varepsilon - \frac{1}{2}}, \quad (\text{D5})$$

and the expansion for small ε

$$\Gamma(1 + \varepsilon) = 1 - \gamma\varepsilon + O(\varepsilon^2), \quad \Gamma\left(\varepsilon + \frac{1}{2}\right) = \Gamma\left(\frac{1}{2}\right) - \varepsilon\Gamma\left(\frac{1}{2}\right)(\gamma + 2\ln 2) + O(\varepsilon^2)$$

$$x^\varepsilon = 1 + \varepsilon \ln x + O(\varepsilon^2), \quad (\text{D6})$$

where γ is the Euler's constant, we find

$$\rho(\varepsilon) = -\frac{U^2(x)}{16} \left[\frac{1}{\varepsilon} + \ln\left(\frac{\mu^2}{U(x)}\right) + 2\ln 2 - \frac{1}{2} \right]. \quad (\text{D7})$$

2. I_- computation

If we define $t = \omega/\sqrt{U(x)}$, the integral I_- in Eq.(D1) becomes

$$\rho(\varepsilon) = \mu^{2\varepsilon} U(x)^{2-\varepsilon} \int_0^{+\infty} dt \frac{t^2}{(t^2 - 1)^{\varepsilon - \frac{1}{2}}} = \frac{1}{2} \mu^{2\varepsilon} U(x)^{2-\varepsilon} B\left(\varepsilon - 2, \frac{3}{2} - \varepsilon\right)$$

$$\frac{1}{2} \mu^{2\varepsilon} U(x)^{2-\varepsilon} \frac{\Gamma\left(\frac{3}{2} - \varepsilon\right) \Gamma(\varepsilon - 2)}{\Gamma\left(-\frac{1}{2}\right)} = -\frac{1}{4\sqrt{\pi}} U(x)^2 \left(\frac{\mu^2}{U(x)}\right)^\varepsilon \Gamma\left(\frac{3}{2} - \varepsilon\right) \Gamma(\varepsilon - 2), \quad (\text{D8})$$

where we have used the following identity involving the beta function

$$\frac{1}{p} B\left(1 - \nu - \frac{\mu}{p}, \nu\right) = \int_1^{+\infty} dt t^{\mu-1} (t^p - 1)^{\nu-1} \quad p > 0, \text{Re } \nu > 0, \text{Re } \mu < p - p \text{Re } \nu \quad (\text{D9})$$

and the reflection formula

$$\Gamma(z) \Gamma(1 - z) = -z \Gamma(-z) \Gamma(z) \quad (\text{D10})$$

From the first of Eqs.(D5) and from the expansion for small ε

$$\Gamma\left(\frac{3}{2} - \varepsilon\right) = \Gamma\left(\frac{3}{2}\right) (1 - \varepsilon(-\gamma - 2\ln 2 + 2)) + O(\varepsilon^2)$$

$$x^\varepsilon = 1 + \varepsilon \ln x + O(\varepsilon^2), \quad (\text{D11})$$

we find

$$\rho(\varepsilon) = -\frac{U^2(x)}{16} \left[\frac{1}{\varepsilon} + \ln\left(\frac{\mu^2}{U(x)}\right) + 2\ln 2 - \frac{1}{2} \right]. \quad (\text{D12})$$

-
- [1] A. G. Riess *et al.*, *Astron. J.* **116**, 1009 (1998);
[2] M. S. Morris and K. S. Thorne, *Am. J. Phys.* **56**, 395 (1988).
[3] M. S. Morris, K. S. Thorne, and U. Yurtsever, *Phys. Rev. Lett.* **61**, 1446 (1988).
[4] F. S. N. Lobo, *Phys. Rev. D* **71** 124022, (2005); gr-qc/0506001. F. S. N. Lobo, *Phys. Rev. D* **71** 084011, (2005); gr-qc/0502099.
[5] P. K. F. Kuhfittig, *Class.Quant.Grav.* **23** 5853, (2006); gr-qc/0608055.

- [6] S. Sushkov, *Phys. Rev. D* **71** 043520, (2005), gr-qc/0502084.
- [7] R. Garattini, *Class.Quant.Grav.* **22** 1105, (2005); gr-qc/0501105.
- [8] R. Garattini and F. Lobo, *Self sustained phantom wormholes in semi-classical gravity* (to be submitted).
- [9] C. J. Fewster, K. D. Olum and M. J. Pfenning, *Averaged null energy condition in spacetimes with boundaries*. gr-qc/0609007.
- [10] D. Hochberg, A. Popov and S. V. Sushkov, *Phys. Rev. Lett.* **78** (1997) 2050, gr-qc/9701064.
- [11] N. R. Khusnutdinov and S. V. Sushkov, *Phys. Rev. D* **65** 084028, (2002), hep-th/0202068.
- [12] M. Visser, *Lorentzian Wormholes* (AIP Press, New York, 1995) 64.
- [13] P. R. Anderson and D. R. Brill, *Phys.Rev. D* **56** (1997) 4824, gr-qc/9610074 .
- [14] R. Arnowitt, S. Deser, and C. W. Misner, in *Gravitation: An Introduction to Current Research* edited by L. Witten (John Wiley & Sons, Inc., New York, 1962).
- [15] R. Garattini, *Phys.Rev. D* **59** 104019, (1999), hep-th/9902006.
- [16] R. Garattini, *Int. J. Mod. Phys. A* **18** (1999) 2905, gr-qc/9805096.
- [17] M. Berger and D. Ebin, *J. Diff. Geom.* **3**, 379 (1969).
- [18] J. W. York Jr., *J. Math. Phys.*, **14**, 4 (1973), *Ann. Inst. Henri Poincaré A* **21** (1974) 319.
- [19] T. Regge and J. A. Wheeler, *Phys. Rev.* **108**, 1063 (1957).
- [20] M. Bordag, U. Mohideen and V.M. Mostepanenko, *Phys. Rep.* **353**, 1 (2001).
- [21] I.O. Cherednikov, *Acta Physica Slovaca*, **52**, (2002), 221.
- [22] S.W. Hawking and G.F.R. Ellis, *The Large Scale Structure of Spacetime* (Cambridge Univ. Press, 1973).
- [23] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series, and Products* (corrected and enlarged edition), edited by A. Jeffrey (Academic Press, Inc.).
- [24] A. P. Prudnikov, Yu. A. Brychkov and O.I. Marichev, *Integrals and Series, Vol. 3: More Special Functions*, edited by Gordon and Breach Science Publishers, Second Printing 1998.